arXiv:1211.4181v1 [math.NT] 18 Nov 2012

Evaluating L-functions with few known coefficients

David W. Farmer
American Institute of Mathematics
farmer@aimath.org

Nathan C. Ryan
Bucknell University
nathan.ryan@bucknell.edu

November 20, 2012

Abstract

We address the problem of evaluating an L-function when only a small number of its Dirichlet coefficients are known. We use the approximate functional equation in a new way and find that is possible to evaluate the L-function more precisely than one would expect from the standard approach. The method, however, requires considerably more computational effort to achieve a given accuracy than would be needed if more Dirichlet coefficients were available.

1 Introduction

L-functions are central to much of contemporary number theory. Two celebrated conjectures, the Riemann Hypothesis and the Birch and Swinnerton-Dyer Conjecture are about values of L-functions and were discovered as a result of the explicit computation of the Riemann zeta function and the Hasse-Weil L-function associated to an elliptic curve, respectively. These L-functions are, respectively, of degree one and degree two and it is interesting to verify analogous conjectures about special values and zeros for higher degree L-functions. Conjectures such as Böcherer's Conjecture [5, 15] and the Bloch-Kato Conjecture [3] are about the central values of degree four and degree three L-functions, and the Grand Riemann Hypothesis asserts that all nontrivial zeros of a "nice" L-function, of any degree, lie along the critical line.

In addition to these conjectures, there are many other conjectures for the statistical behavior of L-functions, arising from random matrix theory and number theoretic heuristics [10, 7, 6]. One of the main reasons those conjectures are believable is that large-scale calculations of the value distribution and the zeros of L-functions yield data that support those conjectures.

The L-functions we consider here are associated to Siegel modular forms. Our examples will use the first non-lift Siegel modular form on $\mathrm{S}p(4,\mathbb{Z})$. The form has weight 20 and is usually denoted Υ_{20}. Background information about Siegel modular forms can be found in [2, 11, 16]. For this paper the relevant information is that a Siegel modular form is acted on by Hecke operators $T(n)$, which have eigenvalues $\lambda(n)$. It is the eigenvalues $\lambda(p)$ and $\lambda(p^2)$ for p prime, which are used to define the L-functions associated to the modular form. For Υ_{20} the eigenvalues $\lambda(p)$ have been computed for $p \leq 997$, and the eigenvalues $\lambda(p^2)$ for $p \leq 79$ [12]. This data is available at [17].

Associated to a Siegel modular form F is a sequence of L-functions, of degrees 4, 5, 10, 14, 16, etc. There is an L-function $L(s, F, \rho)$ of degree n for each n dimensional representation ρ of the dual group of PGSp(4), namely $\mathrm{Sp}(4,\mathbb{C})$. The degree 4, 5, and 10 L-functions are called, respectively, the spinor, standard, and adjoint, and are denoted $L(s, F, \mathrm{spin})$, $L(s, F, \mathrm{stan})$, and $L(s, F, \mathrm{adj})$. Proposition 2.1, taken from [9], summarizes the properties of those L-functions for a weight k Siegel modular form on $\mathrm{S}p(4,\mathbb{Z})$.

The degree 4 and 5 L-functions were shown by Andrianov [1] and Böcherer [4] to have an analytic continuation and satisfy a functional equation. The degree 10 L-function was only recently shown by Pitale, Saha, and Schmidt [13] to have an analytic continuation and satisfy a functional equation. Those properties for the L-functions of degree 14 and above are still conjectural.

1.1 Evaluating L-functions

We are concerned with numerically evaluating L-functions. The standard tool, which is used in popular computational packages [14, 8] is the approximate functional equation. See Proposition 3.1 for the precise formulation.

There are two main difficulties in evaluating high degree L-functions. The first is that if the L-function $L(s)$ has degree d, evaluating $L(\frac{1}{2} + it)$ using the approximate functional equation requires $\gg (1 + |t|)^{d/4}$ Dirichlet series coefficients. Here the implied constant depends on the L-function and the desired precision in the answer. For example, estimating the implied constant from the calculations in Section 4, to find the first 1 million zeros of $L(s, \Upsilon_{20}, \mathrm{adj})$, the degree 10 L-function associated to Υ_{20}, would require using the approximate functional equation with around 1 trillion Dirichlet series coefficients. Note that 1 million zeros is not even a large sample; for example it is probably not sufficient for testing various conjectures about the lower-order terms in the distribution of zeros.

The second difficulty is that current methods are incapable of producing a large number of Dirichlet coefficients of the standard and adjoint L-functions of a Siegel modular form. The Fourier coefficients of index up to 3000000 have been computed for Υ_{20} [12]. These Fourier coefficients are used to compute the Hecke eigenvalues. Examination of formulas on pages 387 of [16] shows that to find the eigenvalue $\lambda(n)$ of $T(n)$, for $n = p^2$, requires the Fourier coefficients of index up to $n^2 = p^4$.

It gets worse. By (2.2) and (2.4), the pth Dirichlet coefficient of the standard or adjoint L-function requires both $\lambda(p)$ and $\lambda(p^2)$. Thus, to determine the first n Dirichlet coefficients of those L-functions requires Fourier coefficients of the Siegel modular form of index up to approximately n^4. The extensive calculations in [12] are not even sufficient to determine the 83rd Dirichlet coefficient of the standard or adjoint L-functions of Υ_{20}.

Of course, one could determine more Dirichlet coefficients by first finding more Fourier coefficients of the cusp form. But the n^4 relationship makes this quite expensive, so with current methods it is not feasible to determine many more Dirichlet coefficients than currently known. It is possible that new methods will be developed to determine the Hecke eigenvalues without extensive computation. The real problem will still remain: how to compute high-degree L-functions without requiring an enormous number of Dirichlet coefficients.

That brings us to the theme of this paper: how accurately can one compute an L-function given a limited number of Dirichlet coefficients. As the above discussion indicates, this is a practical problem and there are many L-functions for which it is not currently possible to determine a reasonably large number of Dirichlet coefficients.

As we describe in this paper, even without knowing many Dirichlet coefficients, we were able to evaluate the L-functions to surprisingly high accuracy (surprising to us, anyway). In fact, we were not able to establish that there is an absolute limit to the accuracy one can obtain from only a limited number of coefficients. However, our method is computationally expensive – much more expensive than evaluating the L-function in a straightforward way if more coefficients were available. If one could find an efficient way to determine the unknown parameters in our method, that could make it possible to quickly evaluate high-degree L-functions. See Section 6 for a discussion.

In the next section we describe the L-functions we consider here, and in Section 3 we recall the approximate functional equation and how it is used to compute an L-function. In Section 4 we state the underlying problem and then describe our experiments evaluating $L(s, \Upsilon_{20}, \text{stan})$ and $L(s, \Upsilon_{20}, \text{adj})$. In Section 5 we describe a second method that performs a little better than the method in Section 4 and show how this second method can be used to approximate unknown Dirichlet series coefficients.

2 The L-functions

The L-functions associated to a Siegel modular form are most conveniently described as Euler products. The local factors of the Euler product can be expressed in terms of the Hecke eigenvalues $\lambda(p)$ and $\lambda(p^2)$, but it is more convenient to express them in terms of the Satake parameters, $\alpha_{0,p}$, $\alpha_{1,p}$, and

$\alpha_{2,p}$, given by

$$\begin{aligned} p^{2k-3} &= \alpha_0^2\alpha_1\alpha_2 \\ A &= \alpha_0^2\alpha_1^2\alpha_2 + \alpha_0^2\alpha_1\alpha_2^2 + \alpha_0^2\alpha_1 + \alpha_0^2\alpha_2 \\ B &= \alpha_0^2\alpha_1^2\alpha_2^2 + \alpha_0^2\alpha_1^2 + \alpha_0^2\alpha_2^2 + \alpha_0^2, \end{aligned} \tag{2.1}$$

where

$$\begin{aligned} \lambda(p)^2 &= 4p^{2k-3} + 2A + B \\ \lambda(p^2) &= (2-1/p)p^{2k-3} + A + B. \end{aligned} \tag{2.2}$$

We suppress the p on the Satake parameters when clear from context.

We rescale the Satake parameters to have the so-called "analytic" normalization $|\alpha_j| = 1$, $\alpha_0^2\alpha_1\alpha_2 = 1$, which is possible if we assume the Ramanujan bound on the Hecke eigenvalues. This corresponds to a simple change of variables in the L-functions, so that all our L-functions satisfy a functional equation in the standard form $s \leftrightarrow 1-s$.

As an error check for the reader who may wish to extend our calculations, for Υ_{20} we have $\lambda(2) = -840960$, $\lambda(4) = 248256200704$, and the Satake parameters at 2 are approximately

$$\begin{aligned} &\{\alpha_0, \alpha_1, \alpha_2\} = \\ &\qquad \{-0.901413 + 0.43296i, 0.630904 - 0.775861i, -0.211226 + 0.977437i\}. \end{aligned} \tag{2.3}$$

Here and throughout this paper, decimal values are truncations of the true values.

Proposition 2.1 *Suppose $F \in M_k(\mathrm{Sp}(4,\mathbb{Z}))$ is a Hecke eigenform. Let $\alpha_{0,p}$, $\alpha_{1,p}$, $\alpha_{2,p}$ be the Satake parameters of F for the prime p, where we suppress the dependence on p in the formulas below. For $\rho \in \{\mathrm{spin}, \mathrm{stan}, \mathrm{adj}\}$ we have the L-functions $L(s,F,\rho) := \prod_{p\ prime} Q_p(p^{-s},F,\rho)^{-1}$ where*

$$\begin{aligned} Q_p(X,F,\mathrm{spin}) &:= (1-\alpha_0X)(1-\alpha_0\alpha_1X)(1-\alpha_0\alpha_2X)(1-\alpha_0\alpha_1\alpha_2X), \\ Q_p(X,F,\mathrm{stan}) &:= (1-X)(1-\alpha_1X)(1-\alpha_1^{-1}X)(1-\alpha_2X)(1-\alpha_2^{-1}X), \\ Q_p(X,F,\mathrm{adj}) &:= (1-X)^2(1-\alpha_1X)(1-\alpha_1^{-1}X)(1-\alpha_2X)(1-\alpha_2^{-1}X) \\ &\qquad (1-\alpha_1\alpha_2X)(1-\alpha_1^{-1}\alpha_2X)(1-\alpha_1\alpha_2^{-1}X)(1-\alpha_1^{-1}\alpha_2^{-1}X), \end{aligned} \tag{2.4}$$

give the L-series of, respectively, the spinor, standard, and adjoint L-functions.

These L-functions satisfy the functional equations:

$$\begin{aligned}
\Lambda(s,F,\mathrm{spin}) &:= \Gamma_{\mathbb{C}}(s+\tfrac{1}{2})\Gamma_{\mathbb{C}}(s+k-\tfrac{3}{2})L(s,F,\mathrm{spin})\\
&= (-1)^k\Lambda(1-s,F,\mathrm{spin}),\\
\Lambda(s,F,\mathrm{stan}) &:= \Gamma_{\mathbb{R}}(s)\Gamma_{\mathbb{C}}(s+k-2)\Gamma_{\mathbb{C}}(s+k-1)L(s,F,\mathrm{stan})\\
&= \Lambda(1-s,F,\mathrm{stan}),\\
\Lambda(s,F,\mathrm{adj}) &:= \Gamma_{\mathbb{R}}(s+1)^2\Gamma_{\mathbb{C}}(s+1)\\
&\quad\times \Gamma_{\mathbb{C}}(s+k-2)\Gamma_{\mathbb{C}}(s+k-1)\Gamma_{\mathbb{C}}(s+2k-3)L(s,F,\mathrm{adj})\\
&= \Lambda(1-s,F,\mathrm{adj}). \qquad (2.5)
\end{aligned}$$

In (2.5), we use the normalized Γ-functions

$$\Gamma_{\mathbb{R}}(s) := \pi^{-s/2}\Gamma(s) \qquad \text{and} \qquad \Gamma_{\mathbb{C}}(s) := 2(2\pi)^{-s}\Gamma(s).$$

The *degree* of an L-function is $r+2c$ where r and c are the number of $\Gamma_{\mathbb{R}}$ and $\Gamma_{\mathbb{C}}$ factors in the functional equation, respectively. The spin, standard, and adjoint L-functions described above are of degree 4, 5, and 10. The Ramanujan bound for a degree d L-function with Dirichlet series $\sum_{n\geq 1} b_n n^{-s}$ is given by:

$$|b_{p^j}| \leq \binom{d+j-1}{j}. \qquad (2.6)$$

Note that this is equivalent to the assertion that the Satake parameters satisfy $|\alpha_{j,p}| \leq 1$.

3 The approximate functional equation

In this section we describe the approximate functional equation, which is the primary tool used to evaluate L-functions. The approximate functional equation involves a test function which can be chosen with some freedom. This will play a key role in our calculations.

3.1 Smoothed approximate functional equations

The material in this section is taken directly from Section 3.2 of [14].

Let

$$L(s) = \sum_{n=1}^{\infty} \frac{b_n}{n^s} \qquad (3.1)$$

be a Dirichlet series that converges absolutely in a half plane, $\Re(s) > \sigma_1$.

Let

$$\Lambda(s) = Q^s \left(\prod_{j=1}^{a} \Gamma(\kappa_j s + \lambda_j)\right) L(s), \qquad (3.2)$$

with $Q, \kappa_j \in \mathbb{R}^+$, $\Re(\lambda_j) \geq 0$, and assume that:

1. $\Lambda(s)$ has a meromorphic continuation to all of $\mathbb{C}$ with simple poles at $s_1, \ldots, s_\ell$ and corresponding residues $r_1, \ldots, r_\ell$.
2. $\Lambda(s) = \varepsilon \overline{\Lambda(1-\overline{s})}$ for some $\varepsilon \in \mathbb{C}$, $|\varepsilon| = 1$.
3. For any $\sigma_2 \leq \sigma_3$, $L(\sigma + it) = O(\exp t^A)$ for some $A > 0$, as $|t| \to \infty$, $\sigma_2 \leq \sigma \leq \sigma_3$, with A and the constant in the 'Oh' notation depending on σ_2 and σ_3.

Note that (3.2) expresses the functional equation in more general terms than (2.5), but it is a simple matter to unfold the definition of $\Gamma_\mathbb{R}$ and $\Gamma_\mathbb{C}$.

To obtain a smoothed approximate functional equation with desirable properties, Rubinstein [14] introduces an auxiliary function. Let $g : \mathbb{C} \to \mathbb{C}$ be an entire function that, for fixed s, satisfies

$$\left|\Lambda(z+s)g(z+s)z^{-1}\right| \to 0 \tag{3.3}$$

as $|\Im z| \to \infty$, in vertical strips, $-x_0 \leq \Re z \leq x_0$. The smoothed approximate functional equation has the following form.

Theorem 3.1 *For $s \notin \{s_1, \ldots, s_\ell\}$, and $L(s)$, $g(s)$ as above,*

$$\Lambda(s)g(s) = \sum_{k=1}^{\ell} \frac{r_k g(s_k)}{s - s_k} + Q^s \sum_{n=1}^{\infty} \frac{b_n}{n^s} f_1(s,n) + \varepsilon Q^{1-s} \sum_{n=1}^{\infty} \frac{\overline{b_n}}{n^{1-s}} f_2(1-s,n) \tag{3.4}$$

where

$$\begin{aligned} f_1(s,n) &:= \frac{1}{2\pi i} \int_{\nu - i\infty}^{\nu + i\infty} \prod_{j=1}^{a} \Gamma(\kappa_j(z+s) + \lambda_j) z^{-1} g(s+z)(Q/n)^z dz \\ f_2(1-s,n) &:= \frac{1}{2\pi i} \int_{\nu - i\infty}^{\nu + i\infty} \prod_{j=1}^{a} \Gamma(\kappa_j(z+1-s) + \overline{\lambda_j}) z^{-1} g(s-z)(Q/n)^z dz \end{aligned} \tag{3.5}$$

with $\nu > \max\{0, -\Re(\lambda_1/\kappa_1 + s), \ldots, -\Re(\lambda_a/\kappa_a + s)\}$.

In our examples $L(s)$ continues to an entire function, so the first sum in (3.4) does not appear. For fixed $Q, \kappa, \lambda, \varepsilon$, and sequence b_n, and $g(s)$ as described below, the right side of (3.4) can be evaluated to high precision.

A reasonable choice for the weight function is $g(s) = e^{ibs+cs^2}$, which by Stirling's formula satisfies (3.3) if $c > 0$, or if $c = 0$ and $|b| < \pi d/4$, where d is the degree of the L-function. Rubinstein [14] uses such a weight function with b chosen to balance the size of the terms in the approximate functional equation, minimizing the loss in precision in the calculation. In this paper we exploit the fact that there are many choices of weight function, and so there are many ways to evaluate the L-function. We combine those calculations to extract as much information as possible from the known Dirichlet coefficients. This idea is described in the next section.

In the computations we carry out below, we find it more convenient to use the Hardy Z-function in our computations instead of the L-function itself. The function Z associated to an L-function L is defined by the properties: $Z(\frac{1}{2}+it)$ is a smooth function which is real if t is real, and $|Z(\frac{1}{2}+it)| = |L(\frac{1}{2}+it)|$.

4 Exploiting the test function in the approximate functional equation

If we let $g(s) = 1$ and $s = \frac{1}{2} + 10i$ in the approximate functional equation (3.4) for the standard (degree 5) L-function of Υ_{20}, we get

$$\begin{aligned} Z(\tfrac{1}{2}+10i, \Upsilon_{20}, \text{stan}) = &-1835.424 - 395.011\, b_2 + 1012.179\, b_3 + 1906.603\, b_4 + \\ &+ 2226.503\, b_5 + \cdots + 6.840 \times 10^{-9}\, b_{82} + \\ &+ 5.132 \times 10^{-9}\, b_{83} + \cdots + 3.205 \times 10^{-16}\, b_{149} \\ &+ 2.564 \times 10^{-16}\, b_{150} + \cdots . \end{aligned} \tag{4.1}$$

If instead we let $g(s) = e^{-3is/2}$ and keep $s = \frac{1}{2} + 10i$ then we have

$$\begin{aligned} Z(\tfrac{1}{2}+10i, \Upsilon_{20}, \text{stan}) = &\, 1.66549 + 1.39643\, b_2 - 0.658439\, b_3 + 0.726149\, b_4 + \\ &- 0.88227\, b_5 + \cdots + 1.532 \times 10^{-8}\, b_{82} \\ &+ 1.271 \times 10^{-8}\, b_{83} + \cdots + 3.514 \times 10^{-14}\, b_{149} \\ &+ 2.309 \times 10^{-14}\, b_{150} + \cdots . \end{aligned} \tag{4.2}$$

Note that neither of the above expressions appears optimal: the first involves large coefficients, which will lead to a loss of precision. In the second, the terms do not decrease as quickly, so one must use more coefficients to achieve a given accuracy. An observation that we exploit is the fact that the above are just two among a large number of expressions for the value of the L-function at $\frac{1}{2} + 10i$.

Recall that by [12, 17] we know the Satake parameters of Υ_{20} for all $p \leq 79$. Therefore we recognize two types of terms in the approximate functional equation, as illustrated in the above examples. There are terms for which we know the Dirichlet coefficients, such as b_5, b_{82}, and b_{150}. And there are terms with an unknown Dirichlet coefficient, such as b_{83} or b_{149}. Actually, there is a third type of term, such as $b_{166} = b_2 b_{83}$ which is "partially unknown". We can estimate the unknown terms by applying the Ramanujan bound to the Dirichlet coefficient, and evaluate everything else precisely. Thus, once we choose a test function, we can evaluate an L-function at a given point as

$$Z(s) = \text{calculated\_value}(s) \pm \text{error\_estimate}(s), \tag{4.3}$$

where both the calculated value and the error estimate are functions of the test function and the set of known Dirichlet coefficients. For later use, we write

$$\text{error\_estimate}(s) = \sum_{n:\ b_n \,\text{unknown}} \delta_n(g,s) b_n \tag{4.4}$$

where the $\delta_n(g,s)$ is the coefficient of b_n in (3.4). The product of $\delta_n(g,s)$ and the Ramanujan bound of the L-function for the coefficient is an upper bound for the error contributed to computation by the unknown coefficient b_n.

Using the known b_n and applying the Ramanujan bound (2.6) to (4.1) we get

$$Z(\tfrac{1}{2}+10i,\Upsilon_{20},\mathrm{stan}) = 3.0393070838088 \pm 3.12\times 10^{-8}. \tag{4.5}$$

And for (4.2) we get

$$Z(\tfrac{1}{2}+10i,\Upsilon_{20},\mathrm{stan}) = 3.0393070808076 \pm 7.10\times 10^{-8}. \tag{4.6}$$

In Figure 1 we show the calculated value and error estimate for $Z(\frac{1}{2}+10i,\Upsilon_{20},\mathrm{stan})$ when evaluated with test functions of the form $g(s)=e^{-i\beta s}$.



Figure 1: The solid line is the calculated value and the dashed line is the error estimate in computing $Z(\frac{1}{2}+10i,\Upsilon_{20},\mathrm{stan})$ using the available Dirichlet coefficients with the weight function $g(s)=e^{-i\beta s}$ where β is given along the horizontal axis. The vertical axis is $\log_{10}$ of (the absolute value of) the actual value. For quite a wide range of test functions, the value of the Z-function at $\frac{1}{2}+10i$ is determined with some accuracy, achieving around 10 decimal digits of accuracy with the optimal choice of β.

As Figure 1 shows, there is a wide range of β for which it is possible to determine $\Lambda(\frac{1}{2}+10i,\Upsilon_{20},\mathrm{stan})$ with some accuracy. With the optimal choice of β the error estimate is approximately 4×10^{-10}.

Figure 2 shows the calculated value and error estimate for the adjoint (degree 10) L-function $Z(\frac{1}{2}+5i,\Upsilon_{20},\mathrm{adj})$ when evaluated with test functions of the form $g(s)=e^{-i\beta s+\frac{1}{500}(s-5i)^2}$.

As Figure 2 shows, every test function of the given form leads to an error which is larger than the calculated value. Thus, we can determine that $|Z(\frac{1}{2}+5i,\Upsilon_{20},\mathrm{adj})|<0.25$, but with individual test functions of the given form we cannot even determine if the actual value is positive or negative.



Figure 2: The analogue of Figure 1 for $Z(\frac{1}{2}+5i, \Upsilon_{20}, \mathrm{adj})$ with the weight function $g(s) = e^{-i\beta s + \frac{1}{500}(s-5i)^2}$ where β is given along the horizontal axis.

We now introduce a new idea for increasing the accuracy of these calculations.

4.1 Optimizing the test function

In Figure 1 we see that there are many values of the parameters in the test function which give reasonable results. If there is some degree of independence in the errors, then there is hope for obtaining a smaller error by combining the results of those separate calculations. Write $Z(s, \Upsilon_{20}, \mathrm{stan}, \beta)$ for the output of the approximate functional equation with weight function $g(s) = e^{-i\beta s}$. Consider

$$Z(s, \Upsilon_{20}, \mathrm{stan}) = \sum_{j=1}^{J} c_{\beta_j} Z(s, \Upsilon_{20}, \mathrm{stan}, \beta_j) \tag{4.7}$$

where $\sum c_{\beta_j} = 1$. We make the specific choices

$$\begin{aligned} s &= \tfrac{1}{2} + 10i \\ (\beta_1, \beta_2, \beta_3, \beta_4, \beta_5) &= \left(\tfrac{1}{10}, \tfrac{1}{5}, \tfrac{3}{10}, \tfrac{2}{5}, \tfrac{1}{2}\right) \\ (c_{\beta_1}, c_{\beta_2}, c_{\beta_3}, c_{\beta_4}, c_{\beta_5}) &= (0.03150, 0.18061, 0.36563, 0.31421, 0.10801). \end{aligned} \tag{4.8}$$

Recall that all decimal numbers are truncations of the actual values; one requires much higher precision than the displayed numbers in order to obtain the answers below. With the choice in (4.8), after substituting the known Dirichlet coefficients and then using the Ramanujan bound, we find

$$\begin{aligned}
&Z(s,\Upsilon_{20},\mathrm{stan}) = \\
&\quad 3.0393070864895278277801 + 2.688\cdot 10^{-19} b_{83} + \cdots - 1.147\cdot 10^{-16} b_{107} + \\
&\quad + \cdots - 5.291\cdot 10^{-18} b_{137} + \cdots + 1.216\cdot 10^{-23} b_{199} + \cdots \\
&= 3.039307086489527827 \pm 4.737\cdot 10^{-15}.
\end{aligned} \tag{4.9}$$

Thus, by averaging only 5 evaluations of the L-function, the error decreased by a factor of almost 10^{-5}.

The weights c_{β_j} in (4.8) were determined by finding the least-squares fit to

$$\sum_{n:\ b_n\ \text{unknown}} Ram(b_n)^2 \Big(\sum_j c_{\beta_j} \delta(\beta_j, \tfrac{1}{2}+10i)\Big)^2 = 0, \tag{4.10}$$

where $Ram(b_n)$ is the Ramanujan bound (2.6) for b_n, subject to $\sum c_{\beta_j} = 1$. Note that in our actual examples the vast majority of unknown coefficients have prime index, so the $Ram(b_n)$ weighting is not important, but we include it for completeness.

The error estimate in (4.9) is an L^1 estimate, not the L^2 estimate as shown in (4.10), so actually it is possible to choose slightly better weights that those used in our example. In Section 5 we show how to obtain the optimal result that can arise from combining different evaluations of the L-function, but for now we merely wish to illustrate the seemingly surprising fact that appropriately combining several evaluations can vastly decrease the error.

4.2 Results

In Figure 3 we plot the error obtained by combining varying numbers of weight functions, where we evaluate $Z(\frac{1}{2}+10i,\Upsilon_{20},\mathrm{stan})$ with a weight function $g(s) = e^{i\beta s}$ with $\beta = j/10$ with $-10 \le j \le 25$. The horizontal axis is the number of terms averaged, where we start with $\beta = \frac{1}{2}$ and first use those β which are closest to $\frac{1}{2}$. The vertical axis is the error estimate on a $\log_{10}$ scale. The lowest point on the graph, when we average all 36 evaluations, corresponds to

$$\begin{aligned}
&Z(\tfrac{1}{2}+10i,\Upsilon_{20},\mathrm{stan}) = \\
&\qquad 3.03930708648952848108246032844222250910 \pm 2.79\times 10^{-35}.
\end{aligned} \tag{4.11}$$

It is not clear from Figure 3 whether one would expect to obtain an arbitrarily small error by combining sufficiently many test functions in the approximate functional equations. This is discussed in Section 6.

We briefly describe the corresponding calculations for $Z(\frac{1}{2}+5i,\Upsilon_{20},\mathrm{adj})$. Recall that, as shown in Figure 2, with a single test function of the standard form we are not able to determine whether that value is positive or negative. Now we combine five evaluations in the analogous way:

$$Z(s,\Upsilon_{20},\mathrm{adj}) = \sum_{j=1}^{J} c_{\beta_j} Z(s,\Upsilon_{20},\mathrm{adj},\beta_j) \tag{4.12}$$



Figure 3: The error obtained from a least-squares minimization of the error for combining n evaluations of $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan})$ using the weight functions $g(s) = e^{-i\beta s}$ with $\beta = j/10$. The horizontal axis is n and the vertical axis is $\log_{10}$ of the resulting error estimate.

where $\sum c_{\beta_j} = 1$. Here the weight function is $g(s) = e^{-i\beta s + (s-5i)^2/500}$.

We make the specific choices

$$\begin{aligned} s &= \tfrac{1}{2} + 5i \\ (\beta_1, \beta_2, \beta_3, \beta_4, \beta_5) &= \left(\tfrac{3}{5}, \tfrac{6}{5}, \tfrac{9}{5}, \tfrac{12}{5}, 3\right) \\ (c_{\beta_1}, c_{\beta_2}, c_{\beta_3}, c_{\beta_4}, c_{\beta_5}) &= (0.035863, 0.33504, 0.47934, 0.13827, 0.01146). \end{aligned} \tag{4.13}$$

The result is

$$Z(\tfrac{1}{2} + 5i, \Upsilon_{20}, \mathrm{adj}) = 0.01556 \pm 0.0049. \tag{4.14}$$

So we have determined that $Z(\frac{1}{2} + 5i, \Upsilon_{20}, \mathrm{adj})$ is positive, but we can only be certain of one significant figure in its decimal expansion. Using this method, the best result we were able to obtain, by averaging 11 evaluations, is

$$Z(\tfrac{1}{2} + 5i, \Upsilon_{20}, \mathrm{adj}) = 0.01558768 \pm 0.00016. \tag{4.15}$$

Averaging more weight function actually makes the result worse. The explanation is simple: since the adjoint L-function has high degree, the error terms δ_n decrease very slowly. The least-squares fit does not properly take into account the contribution of a large number of small terms, so the least-squares fit actually has a large L^1 norm.

In Section 5 we introduce a different method which avoids these shortcomings.

5 Linear programming

In this section we view the evaluation of the L-function as an optimization problem. For example, we can view the equality of the expressions in (4.1) and (4.2) as a *constraint* on the value of the L-function. Thus, the same calculations which were used as input for the least-squares method described in the previous section can also be used as input to a linear programming problem.

We set up the linear programming problem as follows. Let $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan}, g_j)$ denote the evaluation of $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan})$ using the weight function g_j in the approximate functional equation. One evaluation, $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan}, g_1)$, is taken as the objective. The other evaluations are taken in pairs and

$$Z(\tfrac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan}, g_1) - Z(\tfrac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan}, g_j) = 0 \tag{5.1}$$

is interpreted as a constraint. The other constraints come from the Ramanujan bound on the unknown coefficients.

In the above description there are infinitely many unknowns and constraints. We eliminate the unknown coefficients b_n with $n > 1000$ by using the Ramanujan bound, replacing the equality (5.1) by a pair of inequalities. Thus, we have a straightforward linear programming problem, which we use to determine the minimum and maximum possible values of $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan})$.

We implemented this idea using the same set of test functions we used for the least-squares method. The calculations were done in Mathematica, using the built-in `LinearProgramming` function with the `Method -> Simplex` option. In all cases the linear programming method gave better results, but not spectacularly better. Figure 4 shows the ratio of the errors in the results of the two methods, on a $\log_{10}$ scale. For example, when using 30 equations the error from the linear programming approach was approximately 1/10 the error from the least-squares method.

5.1 Computing unknown coefficients

An interesting side-effect of the linear programming approach is that it allows us to obtain information about the unknown coefficients. Instead of treating the value of the L-function as the objective, we can use an unknown coefficient as the objective. Note that all the other constraints in the problem are unchanged. Using the same method as described above we find the following coefficients of the standard L-function of Υ_{20}:

$$\begin{aligned} b_{83} &= 0.4884558312724 \pm 2.4 \times 10^{-12} \\ b_{89} &= 0.10561760640 \pm 2.7 \times 10^{-10} \\ b_{97} &= 0.468135808 \pm 1.5 \times 10^{-7} \end{aligned} \tag{5.2}$$

These results can be checked once more Fourier coefficients of Υ_{20} are computed. By the n^4 argument given in in the Introduction, computing exact values of b_p for $p \leq 97$ will take about twice as much work as it took to compute those for $p \leq 79$.



Figure 4: The ratio of the errors in the linear programming and least-squares methods for evaluating $Z(\frac{1}{2}+10i, \Upsilon_{20}, \mathrm{stan})$ using n equations. The horizontal axis is n and the vertical axis is $\log_{10}$ of the ratio of errors.

6 Conclusions and further questions

We have shown that, at the cost of a lot of computation, one can evaluate an L-function to high precision using only a small number of coefficients. That this is theoretically possible is not surprising: L-functions are very special objects, and the data we have for the L-function considered here (the functional equation and the first several coefficients) presumably specify the L-function uniquely. Thus, in an abstract sense there is no new information in the missing coefficients. But the question remains as to whether our methods accomplish this in practice.

Question 6.1 *Can the method of calculating an L-function by evaluating the approximate functional equation and then averaging to minimize the contributions of the unknown coefficients, determine numerical values of the L-function to arbitrary accuracy?*

Because the approximate functional equation requires a huge number of terms to evaluate a high-degree L-function, it would be significant if the weights we obtained by our methods could be determined without actually calculating all the terms with unknown coefficients.

Problem 6.2 *Devise a method of determining an optimal weight function in the approximate functional equation without first calculating a large number of terms which do not actually contribute to the final answer.*

Implicit in the above problem is the requirement that one knows the calculated value as well as an estimate of the error.

Problem 6.3 *Is there any meaning to the weights determined by the least-squares method?*

The weights which appear in our least-squares method depend on the point at which the L-function is evaluated. It might be helpful to consider a case where there are two L-functions with the same functional equation, such as the spin L-functions of $Sp(4,\mathbb{Z})$ Siegel modular forms of weight $k \geq 22$.

Without progress on these problems, or a completely new method, there is little hope of making extensive computations of high-degree L-functions.

References

[1] A. N. Andrianov. Euler products that correspond to Siegel's modular forms of genus 2. *Uspehi Mat. Nauk*, 29(3 (177)):43–110, 1974.

[2] Anatolij N. Andrianov. *Quadratic forms and Hecke operators*, volume 286 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1987.

[3] Spencer Bloch and Kazuya Kato. L-functions and Tamagawa numbers of motives. In *The Grothendieck Festschrift, Vol. I*, volume 86 of *Progr. Math.*, pages 333–400. Birkhäuser Boston, Boston, MA, 1990.

[4] Siegfried Böcherer. Über die Funktionalgleichung automorpher L-Funktionen zur Siegelschen Modulgruppe. *J. Reine Angew. Math.*, 362:146–168, 1985.

[5] Böcherer, Siegfried. Bemerkungen über die Dirichletreihen von Koecher und Maaß. (Remarks on the Dirichlet series of Koecher and Maaß). Technical report, Math. Gottingensis, Schriftenr. Sonderforschungsbereichs Geom. Anal. 68, 36 S. , 1986.

[6] Brian Conrey, David W. Farmer, and Martin R. Zirnbauer. Autocorrelation of ratios of L-functions. *Commun. Number Theory Phys.*, 2(3):593–636, 2008.

[7] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein, and N. C. Snaith. Integral moments of L-functions. *PLMS*, 91(3):33–104, 2005.

[8] Tim Dokchitser. Computing special values of motivic L-functions. *Experiment. Math.*, 13(2):137–149, 2004.

[9] David W. Farmer, Nathan C. Ryan, and Ralf Schmidt. Testing the functional equation of a high-degree Euler product. *Pac. J. Math.*, 253(2):349,366, 2011.

[10] J. P. Keating and N. C. Snaith. Random matrices and L-functions. *J. Phys. A*, 36(12):2859–2881, 2003. Random matrix theory.

[11] Helmut Klingen. *Introductory lectures on Siegel modular forms*, volume 20 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1990.

[12] Winfried Kohnen and Michael Kuss. Some numerical computations concerning spinor zeta functions in genus 2 at the central point. *Math. Comp.*, 71(240):1597–1607 (electronic), 2002.

[13] Ameya Pitale, Abhishek Saha, and Ralf Schmidt. Transfer of Siegel cusp forms of degree 2. *Preprint*, 2012.

[14] Michael Rubinstein. Computational methods and experiments in analytic number theory. In *Recent perspectives in random matrix theory and number theory*, volume 322 of *London Math. Soc. Lecture Note Ser.*, pages 425–506. Cambridge Univ. Press, Cambridge, 2005.

[15] Nathan C. Ryan and Gonzalo Tornaría. A Böcherer-type conjecture for paramodular forms. *Int. J. Number Theory*, 7(5):1395–1411, 2011.

[16] Nils-Peter Skoruppa. Computations of Siegel modular forms of genus two. *Math. Comp.*, 58(197):381–398, 1992.

[17] Nils-Peter Skoruppa. Data for rational Siegel eigenforms. *Personal website*, 2010.